

ATP-TR-BCRP-M 1.0 VT-nHTS-BCRP-M-E3S 1.0 VT-nHTS-BCRP-M-MXT 1.0 VT-HTS-BCRP-M-MTX 1.0 VT-HTS-BCRP-M-E3S 1.0	 <b>SOLVO</b> Biotechnology	
---	---	--

## Membrane Product Data Sheet [M-CTRL]

**Catalogue number:**   
**Description:**

**Date of production (dd.mmm.yyyy):**   
**Expiry date (dd.mmm.yyyy):**  when stored at  $-80\text{ }^{\circ}\text{C}$

**Packaging:**

**Total volume:**

**Protein concentration:**

**Total protein:**

	Normal Range	Specific activity
<b>Basal vanadate sensitive ATPase activity (nmol Pi/mg/min)</b>	0-5	
<b>ATP dependent E3S transport @1 µM [pmol/mg/min]</b>	<5	
<b>ATP dependent MTX transport @100 µM [pmol/mg/min]</b>	<5	

**Intended use:**

### Methods:

Protein concentrations were determined using the BCA assay. ATPase activity was measured according to the ATPase assay protocol. Vesicular transport was measured according to the corresponding vesicular transport protocol. Protocols are supplied with the membranes.

### Storage and handling:

- Store at  $-80\text{ }^{\circ}\text{C}$
- Thaw membranes in a water bath at  $25\text{ }^{\circ}\text{C}$ , then store on ice until use.
- The baseline vanadate sensitive ATPase activity of the membranes does not decrease significantly after one freeze-thaw cycle. If you are planning to reuse the same vial, minimize the number of freeze-thaw cycles by making smaller aliquots.
- The vesicular structure of the membrane preparation might be destroyed upon freezing and thawing. If you are using a membrane stock that has been thawed and frozen always include membrane validation in your assay (drug free control – see assay protocol for details).

Validated by:

Date: